

$NN(^1S_0)$ pairs in ^3He and in $p^3\text{He}$ backward elastic scattering

Yu.N. Uzikov^{a*} and J. Haidenbauer^b

^aJoint Institute for Nuclear Research, Dubna, Moscow region, Russia, 141980

^b Institut für Kernphysik, Forschungszentrum Jülich, D-52425 Jülich, Germany

It is shown that the shoulder observed in the cross section of $p^3\text{He}$ backward elastic scattering at proton beam energies of 0.4 – 0.7 GeV can be explained by the mechanism of virtual π -meson production with a dominant contribution from the subprocesses $pd^* \rightarrow ^3\text{He} \pi^0$ and $p(pp) \rightarrow ^3\text{He} \pi^+$, where d^* denotes the singlet deuteron and the pp pair is in the 1S_0 state. At high energies, 1 – 3 GeV, the mechanism of np -pair transfer dominates and probes high-momentum $NN(^1S_0)$ correlations in ^3He .

Over the past few years $p^3\text{He}$ backward elastic scattering has been investigated [1, 2] on the basis of the DWBA using a $3N$ bound-state wave function obtained from solving the Faddeev equations for the Reid RSC NN potential. Those studies suggested that this process at beam energies $T_p > 1$ GeV can give unique information about the high momentum component of the ^3He wave function $\varphi^{23}(\mathbf{q}_{23}, \mathbf{p}_1)$, and specifically for high relative momenta, $q_{23} > 0.6$ GeV/c, of the nucleon pair $\{23\}$ in the 1S_0 state and low momenta of the nucleon "spectator" $p_1 < 0.1$ GeV/c. Here φ^{23} is the first Faddeev component of the full wave function of ^3He , $\Psi(1, 2, 3) = \varphi^{23} + \varphi^{31} + \varphi^{12}$. The calculations presented in Refs. [1, 2] demonstrate the dominance of the mechanism of sequential transfer (ST) of the proton-neutron (np) pair (Fig.1a) over a wide range of beam energies, $T_p = 0.1 - 2$ GeV, except for the region of the ST dip at around 0.3 GeV. Other mechanisms of two-nucleon transfer, such as deuteron exchange, non-sequential np transfer [1], and direct pN scattering [3] involve very high internal momenta in the ^3He wave function in q_{23} as well as in p_1 and, as a consequence, give much smaller contributions. As is shown here (see for details Ref. [4]), the region of the ST dip (0.4-0.7 GeV) is dominated by the triangle diagrams of one pion exchange (OPE) with the subprocesses $pd^* \rightarrow ^3\text{He} \pi^0$ and $p(pp) \rightarrow ^3\text{He} \pi^+$ (Fig. 1b), where d^* and pp are the spin-singlet 1S_0 deuteron and diproton, respectively.

In Ref. [2] the deuteron contribution to the cross section of $p^3\text{He}$ scattering within the OPE mechanism is expressed via the experimental cross section of the reaction $pd \rightarrow ^3\text{He} \pi^0$, without elaboration of its concrete mechanism. In order to calculate the contribution of the meson production on the d^* and on the diproton in ^3He , we use the $d^* + p$ and $(pp) + n$ configurations of ^3He calculated in Refs. [5, 6] and adopt the spec-

*Permanent address: Kazakh National University, Institute for Experimental and Theoretical Physics, 480078 Almaty, Kazakhstan; Supported by BMBF (grant KAZ-02/001 and Heisenberg-Landau program)




Figure 1. Right panel: The ST (a) and OPE (b) mechanisms of elastic $p^3\text{He}$ scattering, and the spectator mechanism of $p(\text{NN}) \rightarrow ^3\text{He} \pi$ (c). Left panel: Cms cross section of $p^3\text{He} \rightarrow ^3\text{He} p$ at the scattering angle $\theta_{cm} = 180^\circ$ versus the proton beam energy. Calculations on the basis of the OPE model with the πNN cut-off momentum $\Lambda_\pi = 1.3 \text{ GeV}/c$: 1 – for d in the intermediate state, 2 – $d + d^* + pp$. The result for the nondistorted ST cross section is given by the dotted line 3 (divided by factor of 3 to normalize on the experiment at 1.5 GeV). The dashed curve 4 shows the slope for the counting rule $d\sigma/dt \sim s^{-22}$ [8]. Experimental data are from Refs. [7] ($\circ$), [9] (filled square), [10] (open square), [11] ($\bullet$), and [12] (filled triangle).

tator mechanism for the subprocesses $p(\text{NN})_{s,t} \rightarrow ^3\text{He} \pi$ (Fig.1c) on the singlet (s) and triplet (t) NN pairs. The analysis of the reaction $pd \rightarrow ^3\text{He} \pi^0$ at $T_p < 1 \text{ GeV}$ [6, 13] and the existing data on the inverse reactions of pion capture, i.e. $\pi^+ ^3\text{He} \rightarrow ppp$ and $\pi^- ^3\text{He} \rightarrow pnn$, confirm this mechanism, cf. the discussion in Ref. [4].

Because of the specific spin structure of the OPE amplitude there is no interference between the triplet (M_d) and singlet ($M_{d^*} + M_{pp}$) amplitudes in the spin-averaged sum:

$$\overline{|M_d + M_{d^*} + M_{pp}|^2} = |K|^2 \left\{ \overline{|G_d T_d|^2} + \frac{1}{3} \overline{|G_{d^*}(T_{d^*} + 2 T_{pp})|^2} \right\}. \quad (1)$$

Here T_α ($\alpha = d, d^*$ and pp) is the meson production amplitude $p(\text{NN})_\alpha \rightarrow ^3\text{He} \pi$ and G_α is the structure factor [4]. The factors $\frac{1}{3}$ and 2 in the second term in the curly brackets of Eq. (1) are combinations of isospin coefficients. Furthermore, we assume that the subprocess $pN \rightarrow (\text{NN})_t \pi$ dominates in the upper vertex of the diagram in Fig.1c and that the amplitude $pN \rightarrow (\text{NN})_s \pi$ is negligible. This is true in the Δ -

region [14]. With this approximation there is a relation between the amplitudes of the processes $pd^* \rightarrow {}^3\text{He} \pi^0$ and $p(pp)_s \rightarrow {}^3\text{He} \pi^+$ which follows from isospin invariance, namely $T_{pp} = 2T_{d^*}$. After that one gets the relation $|\overline{T_{d^*} + 2T_{pp}}|^2 = 25|\overline{T_d}|^2$. Therefore the combined contribution of $T_{d^*} + T_{pp}$ is significantly larger than that of the deuteron and can be expressed via the experimental cross section of the reaction $pd \rightarrow {}^3\text{He} \pi^0$ taken here from Ref. [15]. Distortions are included in the factor K .

The results of our calculation are shown in Fig. 1. Evidently, the OPE model with deuteron exchange (curve 1) yields a reasonable description of the energy dependence of the cross section for $T_p = 0.4 - 1.5$ GeV. However, due to distortions, taken into account here in eikonal approximation, the d contribution turns out to be one order of magnitude smaller than the data. The contributions of the singlet deuteron d^* and of the pp pair bring the calculation in qualitative agreement with the data (curve 2).

One can see from curve 3 in Fig. 1 that the ST mechanism, calculated here with the $3N$ wave function [16] resulting from the CD Bonn potential, is significant at beam energies $T_p = 0.9 - 1.5$ GeV and it definitely dominates at low ($T_p < 0.3$ GeV) and high ($T_p > 1.5$ GeV) energies. Since at sufficiently high relative NN momenta the concept of having individual nucleons inside nuclei is expected to fail, we show in Fig.1, for comparison, also the slope of the cross section as it follows from quark counting rules [8] for direct (without baryon exchanges) mechanism. Obviously, this slope differs from the one of the ST and OPE mechanisms.

In conclusion, our calculations suggest that the shoulder in the $p^3\text{He} \rightarrow {}^3\text{He} p$ cross section at 0.4–0.6 GeV is mainly due to the OPE mechanism with the singlet $NN(^1S_0)$ pairs in ${}^3\text{He}$. A measurement of spin observables, planned at 0.2-0.4 GeV [17] could give additional information here.

REFERENCES

1. L.D. Blokhintsev, A.V. Lado, Yu.N. Uzikov, Nucl. Phys. **A597** (1996) 487.
2. Yu.N. Uzikov, Nucl.Phys. **A644** (1998) 321; Phys. Rev. **C 58** (1998) 36.
3. M.J. Paez, R.H. Landau, Phys. Rev. **C 29**(1984) 2267.
4. Yu.N.Uzikov and J.Haidenbauer, Phys. Rev. C 68 (2003) 014001.
5. R. Schiavilla, V.R. Pandharipande, R.B. Wiringa, Nucl. Phys. **A449** (1986) 219.
6. J.-F. Germond, C. Wilkin, J. Phys. G **14** (1988) 181.
7. P. Berthet *et al.*, Phys. Lett. **106B** (1981) 465.
8. S.J. Brodsky and G.R. Farrar, Phys.Rev.Lett. 18 (1973) 1153.
9. H. Langevin-Joliot *et al.*, Nucl. Phys. **A158** (1970) 309.
10. V.I. Komarov *et al.*, Yad. Fiz. **11** (1970) 399.
11. R. Frascaria *et al.*, Phys. Lett. **66B** (1977) 329.
12. L.G. Votta *et al.*, Phys. Rev. **C 10** (1974) 520.
13. J.M. Laget, J.F. Lecolley, Phys. Lett **B194** (1987) 177.
14. Yu.N. Uzikov, C. Wilkin, Phys. Lett. **B545** (2001) 191.
15. P. Berthet *et al.*, Nucl. Phys. **A443** (1985) 589
16. V. Baru, J. Haidenbauer, C. Hanhart, J.A. Niskanen, Eur. Phys. J. **A,16** (2003)437.
17. K. Hatanaka et al., in *Nuclear Many-Body and Medium Effects in Nuclear Interactions and Reactions*, edited by K. Hatanaka et al. (World Scientific, 2003), p. 149.